

On meso-scale approximations for vibrations of membranes with lower-dimensional clusters of inertial inclusions

V.G. Maz'ya^{1,2,3}, A.B. Movchan², M.J. Nieves⁴

¹ Department of Mathematics, Linköping University,
Linköping S-581 83, Sweden

² Department of Mathematical Sciences, University of Liverpool,
Liverpool L69 7ZL, UK

³ RUDN University, 6 Miklukho-Maklay St, Moscow, 117198, Russia

⁴ School of Computing and Mathematics, Keele University,
Staffordshire, ST5 5BG, UK

Abstract

In this paper we consider formal asymptotic algorithms for a class of meso-scale approximations for problems of vibration of elastic membranes containing clusters of small inertial inclusions distributed along contours of pre-defined smooth shapes. Effective transmission conditions have been identified for inertial structured interfaces, and approximations to solutions of eigenvalue problems have been derived for domains containing lower-dimensional clusters of inclusions.

In honour of Professor N.N. Uraltseva

1 Introduction

We address a class of asymptotic approximations for models of vibrations of two-dimensional elastic membranes, containing clusters of small inclusions. These clusters are assumed to be distributed along one-dimensional sets. The approach is based on the method of meso-scale asymptotic approximations [1–4], and the inertia of inclusions has been taken into account. The method of meso-scale asymptotic approximations was first introduced in [1]

for asymptotic problems in domains with large clusters of small inclusions, and it provides an efficient alternative to homogenisation approximations (see, for example, [5]), especially for the cases where clusters are inhomogeneous, non-periodic, and where the size of inclusions is in the meso-scale range compared to the distance between the inclusions. The asymptotic approximations obtained in [1] involve a linear combination of solutions to certain model problems whose coefficients satisfy a linear algebraic system. The solvability of this system was proved under weak geometrical assumptions, and both uniform and energy estimates for the remainder term were derived.

Meso-scale approximations for eigenvalue problems in domains with clusters of many inclusions were analysed in [6]. Fundamental ideas of the method of compound asymptotic expansions [7] in domains with singularly perturbed boundaries have been used.

Although problems of wave scattering in the low frequency regime in solids with many inclusions can be addressed by the method of homogenisation, important features linked to the wave scattering from individual inclusions may require the use of the dynamic Green's functions. In particular, Green's function for the Helmholtz equation in a periodic domain was analysed in [8,9]. Formal procedure was introduced in [10] for isotropic scattering by randomly distributed scatterers separated by a finite distance. Analysis of waves in a plane with semi-infinite arrays of isotropic scatterers is discussed in [11]. Waves governed by the Helmholtz equation in a doubly periodic array with an elementary cell containing several scatterers were analysed in [12], based on the approach of [10], [11] and asymptotics representing singular perturbation leading-order approximations, similar to simplified cases of [7], [2] and [6]. Dispersion of waves analysed in [12] shows dynamic anisotropy linked to the scatterers within the elementary cell.

An example of a cluster of inclusions placed along a curve in a two-dimensional elastic membrane is shown in Fig. 1. Time-harmonic vibrations are considered here, and the inertia of small inclusions, as well as their size, separation and the size of the cluster represent parameters of the multi-scale asymptotic approximation.

When an incident wave of a radian frequency ω is generated by a remote time-harmonic point source placed at \mathbf{y} the approximation of the total field is constructed in the form

$$\mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) \sim G(\mathbf{x}, \mathbf{y}, \omega) + \sum_{j=1}^N \beta_j(\mathbf{y}) U^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|),$$

where $G(\mathbf{x}, \mathbf{y}, \omega)$ is Green's function for the Helmholtz equation, $U^{(j)}$ are special model fields associated with individual small inertial inclusions placed at $\mathbf{O}^{(j)}$, and $\beta^{(j)}$ are the coefficients, found by solving a model algebraic system of equations. The proof of the solvability of this algebraic system is




Figure 1: Cluster of small inclusions placed along a contour. The figure shows the eigenfunction corresponding to the first eigenvalue of the Dirichlet problem for the Laplacian, $\lambda = 0.30816$. The main region is a disk of radius $R = 7$. Small circular rigid inclusions of radii $r = 0.2$ have their centres along the circle of radius 2. The computation is produced in COMSOL.

similar to [1]. Compared to meso-scale two-dimensional clusters, this is a lower-dimensional case where inclusions are distributed along a contour. In addition to the above point-wise asymptotic approximation, the approach discussed here, can also be used to derive an effective transmission problem for the domain containing an inertial structured interface, and its solution provides a homogenisation approximation, which takes into account inertia of small inclusions. In particular, when small inclusions of mass m form a periodic cluster of the overall mass M distributed over a circle Γ of radius $R = 1$, the coefficients β_j can be approximated by

$$\beta_j(\mathbf{y}) = \mathfrak{G}(\mathbf{O}^{(j)}, \mathbf{y}), \quad 1 \leq j \leq N.$$

where the function $\mathfrak{G}(\mathbf{O}^{(j)}, \mathbf{y})$ is the solution of the following transmission problem

$$\mu \Delta_x \mathfrak{G}(\mathbf{x}, \mathbf{y}) + \rho \omega^2 \mathfrak{G}(\mathbf{x}, \mathbf{y}) + \mathfrak{R}(m, M, \omega) \delta(\mathbf{x} - \mathbf{y}) = 0,$$

where μ is the stiffness coefficient, ρ is the mass density, and $\mathfrak{R}(m, M, \omega)$ is the coefficient, which depends on the inertial properties of the cluster. While $\mathfrak{G}(\mathbf{O}^{(j)}, \mathbf{y})$ satisfies the radiation condition at infinity, the transmission conditions across Γ have the form

$$[\mathfrak{G}(\mathbf{x}, \mathbf{y})] = 0, \quad \left[\frac{\partial \mathfrak{G}}{\partial r}(\mathbf{x}, \mathbf{y}) \right] = \frac{M}{2\pi\rho} \mathfrak{R}(m, M, \omega) \mathfrak{G}(\mathbf{x}, \mathbf{y}).$$

For non-resonance forced problems, we also discuss solutions which can be interpreted as fields associated with the time-harmonic scattering from a lower-dimensional cluster of small inertial inclusions. The incident field is produced by a localised source, associated with a forced motion of one of rigid inclusions within the array, and formally the algorithm requires an asymptotic approximation of the harmonic capacitary potential in the low frequency regime. The approximation corresponds to a non-resonance case.

Furthermore, we analyse an asymptotic solutions of an eigenvalue problem for a domain containing a lower-dimensional cluster of small inertial inclusions. When the domain represents a two-dimensional membrane, the cluster of small inclusions is distributed along a simple curve of finite length. The mass density of an individual small inclusion is used as a control parameter for the inertia of the inclusion itself. An example, discussed in the text, gives a comparison between an analytical asymptotic procedure and a FEM simulation produced in COMSOL MultiPhysics shown in Fig. 1, with the analytical logarithmic asymptotic approximation shown in Fig. 2 and producing a remarkably good result.

The structure of the paper is as follows. In Section 2 we introduce the asymptotic model of a one-dimensional cluster of many inertial inclusions placed along a closed simple contour, which forms an inertial structured interface. Section 3 includes analysis of a low-frequency scattering in a finite elastic membrane with a sparse cluster of small movable rigid inclusions. The approximation of the first eigenvalue and the corresponding eigenfunction for a sparse or one-dimensional cluster of inertial inclusions in a finite elastic membrane is discussed in Section 4.

2 Elastic membrane with a one-dimensional cluster of inclusions

The problem considered concerns the scattering and localisation phenomena due to a cluster of N inertial inclusions, $F_\varepsilon^{(j)}$ with masses m_j and centres $\mathbf{O}^{(j)}$, $1 \leq j \leq N$, in a membrane that has density ρ and shear modulus μ . The overall mass of the cluster $M = \sum_{j=1}^N m_j$, is finite. Here we seek the asymptotic approximation of the solution of the scattering problem for a wave initiated by a point source positioned outside the cluster

$$\mu \Delta \mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) + \rho \omega^2 \mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) + \delta(\mathbf{x} - \mathbf{y}) = 0, \quad \mathbf{x} \in \mathbb{R}^2 \setminus \cup_{j=1}^N F_\varepsilon^{(j)}, \quad (1)$$

$$\mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) = C_j, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}, \quad (2)$$

$$\frac{\partial \mathcal{G}}{\partial r_x}(\mathbf{x}, \mathbf{y}, \omega) - ik_0 \mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) = O\left(\frac{1}{r_x^{3/2}}\right), \quad \text{as } r_x \rightarrow \infty, \quad (3)$$

where the position of the point force is given by $\mathbf{y} \in \mathbb{R}^2 \setminus \cup_{j=1}^N \overline{F_\varepsilon^{(j)}}$, with \mathbf{y} being separated by a finite distance from the cluster itself, $r_x = |\mathbf{x}|$ and C_j are constants that are determined in what follows. Here, ω and $k_0 = \sqrt{\rho\omega^2/\mu}$ denote the radian frequency and wavenumber, respectively, of waves in the medium. In addition, as each mass vibrates in the membrane, the linear momentum balance for the mass is taken into account, imposing an additional condition in the form

$$-m_j\omega^2 C_j = \mu \int_{\partial F_\varepsilon^{(j)}} \frac{\partial \mathcal{G}}{\partial n}(\mathbf{x}, \mathbf{y}, \omega) ds_{\mathbf{x}}, \quad (4)$$

for $1 \leq j \leq N$. This problem can be interpreted as that of the time-harmonic scattering of a wave produced by a point source in the presence of a cluster of small inertial inclusions positioned along a simple smooth curve in a plane.

2.1 Model problems

We use the algorithm of the method of meso-scale asymptotic approximations [2] to construct formally an asymptotic approximation of the field $\mathcal{G}(\mathbf{x}, \mathbf{y}, \omega)$. The procedure requires several model problems whose solutions do not depend on the small parameter ε .

2.1.1 Dynamic Green's function

By G we denote the dynamic Green's function for the infinite membrane, that satisfies the equation

$$\mu\Delta G(|\mathbf{x} - \mathbf{y}|, \omega) + \rho\omega^2 G(|\mathbf{x} - \mathbf{y}|, \omega) + \delta(\mathbf{x} - \mathbf{y}) = 0, \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}^2, \quad (5)$$

and the outgoing wave solution has the representation

$$G(|\mathbf{x} - \mathbf{y}|, \omega) = \frac{i}{4\mu} H_0^{(1)}(k_0|\mathbf{x} - \mathbf{y}|), \quad (6)$$

where $k_0 = \sqrt{\rho\omega^2/\mu}$, and $H_0^{(1)}(k_0 r)$ is the Hankel function of the first kind. In particular,

$$G(|\mathbf{x} - \mathbf{y}|, \omega) \sim -\frac{1}{2\pi\mu} \log |k_0|\mathbf{x} - \mathbf{y}||,$$

as $|k_0|\mathbf{x} - \mathbf{y}|| \rightarrow 0$.

2.1.2 Dirichlet problem for the Helmholtz equation in the exterior of a finite inclusion

To allow for the correction of discrepancies on interior boundaries we introduce the problem

$$\mu\Delta U^{(j)}(\mathbf{x}) + \rho\omega^2 U^{(j)}(\mathbf{x}) = 0, \quad \mathbf{x} \in \mathbb{R}^2 \setminus F_\varepsilon^{(j)}, \quad (7)$$

$$U^{(j)}(\mathbf{x}) = 1, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}, \quad (8)$$

where the $U^{(j)}$ also satisfies the radiation condition at infinity:

$$\frac{\partial U^{(j)}}{\partial r}(\mathbf{x}) - ikU^{(j)}(\mathbf{x}) = O\left(\frac{1}{r^{3/2}}\right), \quad \text{as } r \rightarrow \infty. \quad (9)$$

2.2 Meso-scale approximation - formal asymptotics

The leading order approximation for the solution of (1)–(4) is sought in the form

$$\mathcal{G}(\mathbf{x}, \mathbf{y}, \omega) \sim G(|\mathbf{x} - \mathbf{y}|, \omega) + \sum_{j=1}^N \beta_j(\mathbf{y}) U^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|), \quad (10)$$

where the coefficients β_j are to be determined.

2.2.1 The algebraic system

It follows from the boundary conditions (2) that to leading order we have

$$C_k = G(|\mathbf{O}^{(k)} - \mathbf{y}|, \omega) + \beta_k(\mathbf{y}) + \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) U^{(j)}(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|), \quad \mathbf{x} \in \partial F_\varepsilon^{(k)}, \quad (11)$$

with $1 \leq k \leq N$. Using the “equations of motions” (4) for individual inclusions together with (10), we deduce that the constants C_k , $1 \leq k \leq N$, also satisfy the relations

$$\begin{aligned} -\frac{m_k \omega^2}{\mu} C_k &= \int_{\partial F_\varepsilon^{(k)}} \frac{\partial \mathcal{G}(\mathbf{x}, \mathbf{y}, \omega)}{\partial n} ds_x \\ &= \int_{\partial F_\varepsilon^{(k)}} \frac{\partial G(|\mathbf{x} - \mathbf{y}|, \omega)}{\partial n} ds_x + \beta_k(\mathbf{y}) \int_{\partial F_\varepsilon^{(k)}} \frac{\partial U^{(k)}(\mathbf{x} - \mathbf{O}^{(k)})}{\partial n} ds_x \\ &+ \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) \int_{\partial F_\varepsilon^{(k)}} \frac{\partial U^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|)}{\partial n} ds_x, \end{aligned} \quad (12)$$

where \mathbf{n} stands for the unit outward normal on $\partial F_\varepsilon^{(j)}$. Taking into account (5), (7), (8), and when $j \neq k$, using the integration by parts one can represent the boundary integrals in (12) via the integrals over $F_\varepsilon^{(j)}$, and hence the representation (12) takes the form

$$\begin{aligned} m_k \omega^2 C_k &= -\mu \beta_k(\mathbf{y}) \int_{\partial F_\varepsilon^{(k)}} \frac{\partial U^{(k)}(\mathbf{x} - \mathbf{O}^{(k)})}{\partial n} ds_x \\ &+ \rho \omega^2 \int_{F_\varepsilon^{(k)}} \left\{ G(|\mathbf{x} - \mathbf{y}|, \omega) + \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) U^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|) \right\} d\mathbf{x}. \end{aligned} \quad (13)$$

Combining (11) and (13), we derive the algebraic system of equations for the coefficients β_j , $1 \leq j \leq N$, in the asymptotic approximation (10)

$$\begin{aligned}
& m_k \omega^2 G(\mathbf{O}^{(k)}, \mathbf{y}, \omega) - \rho \omega^2 \int_{F_\varepsilon^{(k)}} G(|\mathbf{x} - \mathbf{y}|, \omega) d\mathbf{x} \\
& + \beta_k(\mathbf{y}) \left\{ m_k \omega^2 + \mu \int_{\partial F_\varepsilon^{(k)}} \frac{\partial U^{(k)}(\mathbf{x} - \mathbf{O}^{(k)})}{\partial n} ds_x \right\} \\
& + m_k \omega^2 \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) \left\{ U^{(j)}(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|) - \frac{\rho}{m_k} \int_{F_\varepsilon^{(k)}} U^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|) d\mathbf{x} \right\} = 0.
\end{aligned} \tag{14}$$

The justification of solvability of the above algebraic systems for the coefficients β_j is similar to [1], and we do not discuss it here.

In particular, when inclusions $F_\varepsilon^{(j)}$ are circular with radii $\varepsilon r^{(j)}$ and centres $\mathbf{O}^{(j)}$, where ε is a small non-dimensional parameter, we have

$$U^{(j)}(\mathbf{x} - \mathbf{O}^{(j)}) = \alpha_\varepsilon^{(j)} G(|\mathbf{x} - \mathbf{O}^{(j)}|, \omega), \tag{15}$$

where

$$\alpha_\varepsilon^{(j)} = -\frac{4i\mu}{H_0^{(1)}(k_0 \varepsilon r^{(j)})}.$$

In this case, the algebraic system (14) for the coefficients β_j , $1 \leq j \leq N$ becomes

$$\begin{aligned}
& m_k G(\mathbf{O}^{(k)}, \mathbf{y}, \omega) - \rho \int_{F_\varepsilon^{(k)}} G(\mathbf{x}, \mathbf{y}, \omega) d\mathbf{x} \\
& + \beta_k \left(m_k + \frac{2\pi\mu\varepsilon r^{(k)} \alpha_\varepsilon^{(k)}}{\omega^2} \frac{G(r)}{\partial r} (r) \Big|_{r=\varepsilon r^{(k)}} \right) \\
& + \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j \alpha_\varepsilon^{(j)} \left\{ m_k G^{(j)}(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|, \omega) \right. \\
& \left. - \rho \int_{F_\varepsilon^{(k)}} G^{(j)}(|\mathbf{x} - \mathbf{O}^{(j)}|, \omega) d\mathbf{x} \right\} = 0.
\end{aligned} \tag{16}$$

2.3 Scattering by a cluster of identical inclusions placed along a circular contour

Here, we assume that the circular inclusions $F_\varepsilon^{(j)}$ have the same radii ε and mass m and that their centres are uniformly distributed along the unit circular contour $\Gamma_1 = \{\mathbf{x} : |\mathbf{x}| = 1\}$. Also, let $d = 2\pi/N$, and assume that $\varepsilon < \text{Const } d^{3/2}$. We analyse the case when $N \rightarrow \infty$, while $M = mN$ is fixed, and obtain a problem concerning a membrane with an inertial ring.

When the inclusions $F_\varepsilon^{(k)}$, $1 \leq k \leq N$, are all circular, some of the integrals appearing in (16) can be evaluated explicitly. In particular, when \mathbf{y} is separated by a finite distance from the cluster of the inclusions we have Using the Graf's addition theorem and expanding the Hankel function in a series we have:

$$H_0^{(1)}(k_0|\mathbf{x} - \mathbf{y}|) = \sum_{\nu=-\infty}^{\infty} H_\nu^{(1)}(k_0|\mathbf{y} - \mathbf{O}^{(k)}|)J_\nu(k_0|\mathbf{x} - \mathbf{O}^{(k)}|)e^{i\nu(\pi - \theta_k + \theta_{k,\mathbf{y}})} .$$

where θ_k is the polar angle of \mathbf{x} measured with respect to $\mathbf{O}^{(k)}$ and $\theta_{k,\mathbf{y}}$ is the polar angle of $\mathbf{O}^{(k)}$ measured with respect to \mathbf{y} . Hence, as $\varepsilon \rightarrow 0$,

$$\int_{F_\varepsilon^{(k)}} G(|\mathbf{x} - \mathbf{y}|, \omega) ds_{\mathbf{x}} = \pi\varepsilon^2 G(|\mathbf{O}^{(k)} - \mathbf{y}|, \omega) + O(\varepsilon^3) = O(\varepsilon^2),$$

and for $j \neq k$, it is derived that

$$\begin{aligned} \int_{F_\varepsilon^{(k)}} G(|\mathbf{x} - \mathbf{O}^{(j)}|, \omega) ds_{\mathbf{x}} &= \pi\varepsilon^2 G(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|, \omega) \\ &+ O(\varepsilon^2 d) = O(\varepsilon^2 |\log d|). \end{aligned}$$

We also note that

$$2\pi\varepsilon\mu r^{(k)} \frac{\partial G}{\partial r}(r, \omega) \Big|_{r=\varepsilon r^{(k)}} = -1 + O(\varepsilon^2 |\log \varepsilon|).$$

For the cluster of identical inclusions, the system (16) becomes

$$\begin{aligned} mG(|\mathbf{O}^{(k)} - \mathbf{y}|, \omega) + \beta_k(\mathbf{y}) \left(m - \frac{\alpha_\varepsilon}{\omega^2} \right) \\ + m\alpha_\varepsilon \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) G(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|) = 0. \end{aligned} \tag{17}$$

where $1 \leq k \leq N$, and

$$\alpha_\varepsilon = -\frac{4i\mu}{H_0^{(1)}(k_0\varepsilon)} .$$

Periodicity and inertia of the cluster

Assume that positions of the centres of inclusions $F_\varepsilon^{(j)}$ are

$$\mathbf{O}^{(j)} = (\cos(2\pi(j-1)/N), \sin(2\pi(j-1)/N))^T, \quad 1 \leq j \leq N .$$

and let $\mathfrak{G}(\mathbf{x}, \mathbf{y})$ be a function such that

$$\beta_j(\mathbf{y}) = \mathfrak{G}(\mathbf{O}^{(j)}, \mathbf{y}), \quad 1 \leq j \leq N . \tag{18}$$

Derivation of the transmission conditions for \mathfrak{G} in the auxiliary problem

Equations (17) can be re-written in the form

$$mG(|\mathbf{O}^{(k)} - \mathbf{y}|, \omega) + \beta_k(\mathbf{y}) \left(m - \frac{\alpha_\varepsilon}{\omega^2} \right) + \frac{M\alpha_\varepsilon}{2\pi} \sum_{\substack{j \neq k \\ 1 \leq j \leq N}} \beta_j(\mathbf{y}) G(|\mathbf{O}^{(k)} - \mathbf{O}^{(j)}|) \frac{2\pi}{N} = 0. \quad (19)$$

By considering the sum in (19) as the Riemann sum and taking the limit as $N \rightarrow \infty$, we arrive at

$$m\omega^2 G(|\mathbf{x} - \mathbf{y}|, \omega) + \mathfrak{G}(\mathbf{x}, \mathbf{y})(m\omega^2 - \alpha_\varepsilon) + \frac{M\omega^2\alpha_\varepsilon}{2\pi} \int_0^{2\pi} G(|\mathbf{x} - \boldsymbol{\eta}|, \omega) \mathfrak{G}(\boldsymbol{\eta}, \mathbf{y}) d\theta_\eta = 0. \quad (20)$$

When this equation is extended to \mathbf{x} outside Γ_1 , one can apply the Laplacian in \mathbf{x} , use (5) and derive

$$0 = -m\omega^2 \left(\frac{\rho\omega^2}{\mu} G(|\mathbf{x} - \mathbf{y}|, \omega) + \frac{1}{\mu} \delta(\mathbf{x} - \mathbf{y}) \right) + (m\omega^2 - \alpha_\varepsilon) \Delta_x \mathfrak{G}(\mathbf{x}, \mathbf{y}) - \frac{M\omega^4\alpha_\varepsilon\rho}{2\pi\mu} \int_{\Gamma_1} G(|\mathbf{x} - \boldsymbol{\eta}|, \omega) \mathfrak{G}(\boldsymbol{\eta}, \mathbf{y}) d\theta_\eta. \quad (21)$$

Furthermore, using (20) and (21) we obtain the equation for $\mathfrak{G}(\mathbf{x}, \mathbf{y})$ in the form

$$\mu \Delta_x \mathfrak{G}(\mathbf{x}, \mathbf{y}) + \rho\omega^2 \mathfrak{G}(\mathbf{x}, \mathbf{y}) + \frac{\rho\omega^2}{\alpha_\varepsilon - m\omega^2} \delta(\mathbf{x} - \mathbf{y}) = 0. \quad (22)$$

In addition, we take the normal derivative in (20) and use the following relations when $\mathbf{x} \in \Gamma_1$

$$\lim_{\boldsymbol{\xi} \rightarrow \Gamma_1^\pm} \int_{\Gamma_1} \frac{\partial G}{\partial r} (|\boldsymbol{\xi} - \boldsymbol{\eta}|, \omega) \mathfrak{G}(\boldsymbol{\eta}, \mathbf{y}) ds_\eta = \mp \frac{1}{2} \mathfrak{G}(\mathbf{x}, \mathbf{y}) + \int_{\Gamma_1} \frac{\partial G}{\partial r} (|\mathbf{x} - \boldsymbol{\eta}|, \omega) \mathfrak{G}(\boldsymbol{\eta}, \mathbf{y}) ds_\eta,$$

(see [13]), and obtain the transmission condition across Γ_1

$$[\mathfrak{G}(\mathbf{x}, \mathbf{y})] = 0, \quad \left[\frac{\partial \mathfrak{G}}{\partial r}(\mathbf{x}, \mathbf{y}) \right] = -\frac{M\omega^2}{2\pi(m\omega^2 - \alpha_\varepsilon)} \mathfrak{G}(\mathbf{x}, \mathbf{y}), \quad (23)$$

with $[\mathfrak{G}(\mathbf{x}, \mathbf{y})]$ denoting the jump of $\mathfrak{G}(\mathbf{x}, \mathbf{y})$ across Γ_1 , while the point \mathbf{y} is separated by a finite distance from Γ_1 :

$$[\mathfrak{G}(\mathbf{x}, \mathbf{y})] = \mathfrak{G}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{x} \in \Gamma_1^+} - \mathfrak{G}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{x} \in \Gamma_1^-}.$$

Thus, the function $\mathfrak{G}(\mathbf{x}, \mathbf{y})$ can be defined as a solution of (22), (23), subject to the radiation condition at infinity. This provides an alternative homogenisation approximation (18) for the coefficients d_j in (10), which takes into account the inertial transmission conditions across the structured interface formed by a cluster of small inclusions.

3 Low-frequency scattering in a finite elastic membrane with a sparse cluster of small movable rigid inclusions

Consider a finite elastic membrane $\Omega \subset \mathbb{R}^2$ containing a sparse cluster with a finite number of small rigid inclusions $F_\varepsilon^{(j)}, j = 1, \dots, N$ of zero mass density, containing interior points $\mathbf{O}^{(j)}$. Given the mass density ρ and a time-harmonic vibration of small radian frequency ω , applied to the inclusion $F_\varepsilon^{(1)}$, the amplitude of the out-of-plane displacement satisfies the problem:

$$\mu \Delta u(\mathbf{x}) + \rho \omega^2 u(\mathbf{x}) = 0, \quad \mathbf{x} \in \Omega_N \quad (24)$$

$$u(\mathbf{x}) = 1, \quad \mathbf{x} \in \partial F_\varepsilon^{(1)} \quad (25)$$

$$u(\mathbf{x}) = A_j, \quad 1 < j \leq N, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)} \quad (26)$$

$$u(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial \Omega \quad (27)$$

where

$$\int_{\partial F_\varepsilon^{(j)}} \frac{\partial u}{\partial n} ds = 0, \quad 1 < j \leq N. \quad (28)$$

The constants A_j are to be determined, and it is assumed that small inclusions $F_\varepsilon^{(j)}$ are separated by a finite distance. We introduce a small non-dimensional parameter f , with $\varepsilon < f$, in such a way that $\rho \omega^2 = f \lambda$. Also, we use the normalisation $\mu = 1$, and hence the equation (24) takes the form

$$\Delta u(\mathbf{x}) + f \lambda u(\mathbf{x}) = 0, \quad \mathbf{x} \in \Omega_N. \quad (29)$$

Also, the notation $F^{(j)}$ is used for scaled inclusions, such that $F^{(j)} = \{\mathbf{x} : \varepsilon \mathbf{x} + \mathbf{O}^{(j)} \in F_\varepsilon^{(j)}\}$

3.1 Green's function and the relative capacity potential

We use the result of [2] and employ the *relative capacity potential* $P_\varepsilon^{(1)}(\mathbf{x})$, which satisfies the boundary value problem in $\Omega \setminus \overline{F_\varepsilon^{(1)}}$

$$\Delta P_\varepsilon^{(1)}(\mathbf{x}) = 0, \quad \mathbf{x} \in \Omega \setminus \overline{F_\varepsilon^{(1)}}, \quad (30)$$

$$u(\mathbf{x}) = 1, \quad \mathbf{x} \in \partial F_\varepsilon^{(1)}, \quad (31)$$

$$u(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial \Omega. \quad (32)$$

Also, the notations $G(\mathbf{x}, \mathbf{y})$ and $g(\mathbf{x}, \mathbf{y})$ are used for Green's functions in Ω and $\mathbb{R}^2 \setminus \overline{F^{(1)}}$, respectively. Thus,

$$\Delta G(\mathbf{x}, \mathbf{y}) + \delta(\mathbf{x} - \mathbf{y}) = 0, \quad \mathbf{x}, \mathbf{y} \in \Omega, \quad (33)$$

$$G(\mathbf{x}, \mathbf{y}) = 0, \quad \mathbf{x} \in \partial \Omega, \quad \mathbf{y} \in \Omega. \quad (34)$$

and

$$\Delta g(\boldsymbol{\xi}, \boldsymbol{\eta}) + \delta(\boldsymbol{\xi} - \boldsymbol{\eta}) = 0, \quad \boldsymbol{\xi}, \boldsymbol{\eta} \in \mathbb{R}^2 \setminus \overline{F}^{(1)}, \quad (35)$$

$$g(\boldsymbol{\xi}, \boldsymbol{\eta}) = 0, \quad \boldsymbol{\xi} \in \partial F^{(1)}, \boldsymbol{\eta} \in \mathbb{R}^2 \setminus \overline{F}^{(1)}, \quad (36)$$

$$g(\boldsymbol{\xi}, \boldsymbol{\eta}) \text{ is bounded as } |\boldsymbol{\xi}| \rightarrow \infty \text{ and } \boldsymbol{\eta} \in \mathbb{R}^2 \setminus \overline{F}^{(1)}. \quad (37)$$

According to Lemma 1.4 in [2], the relative capacity potential $P_\varepsilon^{(1)}$ has the asymptotic representation

$$P_\varepsilon^{(1)}(\mathbf{x}) = \frac{-G(\mathbf{x}, \mathbf{O}^{(1)}) + \zeta\left(\frac{\mathbf{x} - \mathbf{O}^{(1)}}{\varepsilon}\right) - \frac{1}{2\pi} \log \frac{|\mathbf{x} - \mathbf{O}^{(1)}|}{\varepsilon r_F}}{\frac{1}{2\pi} \log \frac{\varepsilon r_F}{R_\Omega}} + p_\varepsilon(\mathbf{x}), \quad (38)$$

where $p_\varepsilon(\mathbf{x}) = O(\varepsilon |\log \varepsilon|^{-1})$ uniformly with respect to $\mathbf{x} \in \Omega \setminus \overline{F}_\varepsilon^{(1)}$. In the above formula (38), the notations r_F and R_Ω stand for the inner conformal radius of $F^{(1)}$ with respect to $\mathbf{O}^{(1)}$, and the outer conformal radius of Ω with respect to $\mathbf{O}^{(1)}$, respectively, as discussed in Section 1.2.1 of [2]. The function ζ is defined by

$$\zeta(\boldsymbol{\eta}) = \lim_{|\boldsymbol{\zeta}| \rightarrow \infty} g(\boldsymbol{\zeta}, \boldsymbol{\eta}). \quad (39)$$

In particular, for the case when the inclusion $F_\varepsilon^{(1)}$ is circular of radius εr_F , the formula (38) simplifies as

$$P_\varepsilon^{(1)}(\mathbf{x}) = \beta_1 G(\mathbf{x}, \mathbf{O}^{(1)}) + O(\varepsilon), \quad \beta_1 = -\left(\frac{1}{2\pi} \log \varepsilon r_F + H(\mathbf{O}^{(1)}, \mathbf{O}^{(1)})\right)^{-1}, \quad (40)$$

where the regular part $H(\mathbf{x}, \mathbf{y})$ of Green's function G is a harmonic function defined by

$$H(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi} \log |\mathbf{x} - \mathbf{y}|^{-1} - G(\mathbf{x}, \mathbf{y}). \quad (41)$$

3.2 Formal asymptotic approximation

The asymptotic approximation is sought in the form

$$u(\mathbf{x}) \sim P_\varepsilon^{(1)}(\mathbf{x}) + f u^{(2)}(\mathbf{x}), \quad (42)$$

and the constants A_j , $j = 2, \dots, N$, representing the rigid motion displacements of massless inclusions, are

$$A_j \sim P_\varepsilon^{(1)}(\mathbf{O}^{(j)}) + f A_j^{(2)}, \quad (43)$$

whereas $u^{(2)}$ satisfies the boundary value problem

$$\Delta u^{(2)}(\mathbf{x}) + \lambda P_\varepsilon^{(1)}(\mathbf{x}) = 0 \text{ in } \Omega_N, \quad (44)$$

$$u^{(2)}(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial F_\varepsilon^{(1)}, \quad (45)$$

$$u^{(2)}(\mathbf{x}) = A_j^{(2)}, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}, j = 2, \dots, N, \quad (46)$$

$$u^{(2)}(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial \Omega. \quad (47)$$

By assuming a circular shape of the inclusion $F_\varepsilon^{(1)}$, using formula (40), and introducing an auxiliary problem

$$\Delta V(\mathbf{x}) + \lambda\beta_1 G(\mathbf{x}, \mathbf{O}^{(1)}) = 0 \text{ in } \Omega, \quad (48)$$

$$V(\mathbf{x}) = 0, \mathbf{x} \in \partial\Omega, \quad (49)$$

we deduce

$$V(\mathbf{x}) = \lambda\beta_1 \int_{\Omega} G(\mathbf{z}, \mathbf{x}) G(\mathbf{z}, \mathbf{O}^{(1)}) d\mathbf{z}, \quad (50)$$

and

$$A_j^{(2)} = V(\mathbf{O}^{(j)}) - V(\mathbf{O}^{(1)}) P_\varepsilon^{(1)}(\mathbf{O}^{(j)}) + O(\varepsilon). \quad (51)$$

In this case, the approximation (42) can be rewritten in the form

$$u(\mathbf{x}) = P_\varepsilon^{(1)}(\mathbf{x}) + f(V(\mathbf{x}) - V(\mathbf{O}^{(1)}) P_\varepsilon^{(1)}(\mathbf{x})) + r(x), \quad (52)$$

where the remainder term satisfies the problem

$$\begin{aligned} \Delta r(\mathbf{x}) + f\lambda r(\mathbf{x}) &= f\lambda(P_\varepsilon^{(1)}(\mathbf{x}) - \beta_1 G(\mathbf{x}, \mathbf{O}^{(1)})) \\ &\quad - f^2(V(\mathbf{x}) - V(\mathbf{O}^{(1)}) P_\varepsilon^{(1)}(\mathbf{x})), \\ r(\mathbf{x}) &= 0 \text{ on } \partial\Omega, \\ r(\mathbf{x}) &= -f(V(\mathbf{x}) - V(\mathbf{O}^{(1)})) \text{ on } \partial F_\varepsilon^{(1)}, \\ r(\mathbf{x}) &= A_j - P_\varepsilon^{(1)}(\mathbf{O}^{(j)}) - f(V(\mathbf{O}^{(j)}) - V(\mathbf{O}^{(1)}) P_\varepsilon^{(1)}(\mathbf{O}^{(j)})) \\ &\quad \text{on } \partial F_\varepsilon^{(j)}, j = 2, \dots, N. \end{aligned}$$

For the case when the inclusion $F_\varepsilon^{(1)}$ is circular, taking into account that the small quantity $f\lambda$ is separated from the spectrum, and using the formulae (43), (51) and (52) we deduce

$$r(\mathbf{x}) = O(f\varepsilon). \quad (53)$$

4 The first positive eigenvalue for a lower-dimensional cluster of inertial inclusions in a finite elastic membrane

Section 3 has addressed low frequency non-resonance vibrations for a sparse cluster of massless inclusions (each individual inclusion has zero inertia). In this case, Green's function for the Laplacian is used as one of the model solutions. If the small inclusions have a non-zero inertia the problem requires Green's function for the Helmholtz operator, which takes into account time-harmonic vibrations of the multi-scale system. In this section, we consider the eigenvalue problem for a finite membrane $\Omega \subset \mathbb{R}^2$, and contained in this membrane is a sparse or lower-dimensional cluster of small inertial inclusions

$F_\varepsilon^{(j)}$ of the mass m_j , $1 \leq j \leq N$. In this case, if d is the minimum distance between two neighbouring inclusions within the cluster, we assume that $dN = O(1)$. As before, we use the notation $\Omega_N := \Omega \setminus \cup_{j=1}^N \bar{F}_\varepsilon^{(j)}$. The eigenfunction u_N and the corresponding eigenvalue λ_N are defined as the solution of the problem

$$\Delta u_N(\mathbf{x}) + \lambda_N u_N(\mathbf{x}) = 0, \quad \mathbf{x} \in \Omega_N, \quad (54)$$

$$u_N(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial\Omega \quad (55)$$

$$u_N(\mathbf{x}) = A_j, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}, \quad (56)$$

with

$$-\gamma_j \lambda_N A_j = \int_{\partial F_\varepsilon^{(j)}} \frac{\partial u_N(\mathbf{x})}{\partial n} ds, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}. \quad (57)$$

where $\lambda_N = \rho \omega_N^2 / \mu$ and $\gamma_j = \frac{m_j}{\rho}$, $1 \leq j \leq N$, and n is the unit outward normal with respect to $F_\varepsilon^{(j)}$. The quantities γ_j represent the equivalent inertial area, which would be required for the inclusion of mass density ρ to have the mass m_j . For convenience, we also introduce the normalisation $\mu = 1$ and the notation $\lambda = \rho \omega^2$, where ω is the radian frequency of the time-harmonic vibrations and ρ is the mass density of the membrane. It is also assumed that $\sum_{j=1}^N m_j = M = O(1)$, and thus $\gamma_j = O(M/(N\rho))$.

Here, we are interested in evaluating the first positive eigenvalue, which takes into account the inertia of the small inclusions $F_\varepsilon^{(j)}$.

4.1 Model problem: Green's function for a finite membrane

Assuming a non-resonance regime for λ , we shall use Green's function $G_\Omega(\mathbf{x}, \mathbf{y}, \lambda)$ in Ω , i.e. the solution of the following problem

$$\Delta G_\Omega(\mathbf{x}, \mathbf{y}, \lambda) + \lambda G_\Omega(\mathbf{x}, \mathbf{y}, \lambda) + \delta(\mathbf{x} - \mathbf{y}) = 0, \quad \mathbf{x} \in \Omega, \quad (58)$$

$$G_\Omega(\mathbf{x}, \mathbf{y}, \lambda) = 0, \quad \mathbf{x} \in \partial\Omega, \quad (59)$$

and the regular part R_Ω of G is defined as

$$R_\Omega(\mathbf{x}, \mathbf{y}, \lambda) = -\frac{1}{4} Y_0(\sqrt{\lambda} |\mathbf{x} - \mathbf{y}|) - G_\Omega(\mathbf{x}, \mathbf{y}, \lambda), \quad (60)$$

where $Y_0(\sqrt{\lambda} r)$ is the Bessel function of the second kind, and

$$Y_0(\sqrt{\lambda} r) \sim \frac{2}{\pi} \log(\sqrt{\lambda} r), \quad \text{as } \sqrt{\lambda} r \rightarrow 0.$$

The auxiliary functions $V^{(j)}$ (compare with (7),(8)) are defined by

$$\Delta V^{(j)}(\mathbf{x}, \lambda) + \lambda V^{(j)}(\mathbf{x}, \lambda) = 0, \quad \mathbf{x} \in \Omega \setminus F_\varepsilon^{(j)}, \quad (61)$$

$$V^{(j)}(\mathbf{x}, \lambda) = 1, \quad \mathbf{x} \in \partial F_\varepsilon^{(j)}, \quad (62)$$

$$V^{(j)}(\mathbf{x}, \lambda) = 0, \quad \mathbf{x} \in \partial\Omega. \quad (63)$$

When $F_\varepsilon^{(j)}$ are small circular inclusions of radii εr_j , $V^{(j)}$ are approximated in the form

$$V^{(j)}(\mathbf{x}, \lambda) = \beta_\varepsilon^{(j)} G_\Omega(\mathbf{x}, \mathbf{O}^{(j)}, \lambda) + O(\varepsilon |\log \varepsilon|^{-1}), \quad (64)$$

where

$$\beta_\varepsilon^{(j)} = -\left(\frac{1}{4} Y_0(\sqrt{\lambda} \varepsilon r_j) + R_\Omega(\mathbf{O}^{(j)}, \mathbf{O}^{(j)}, \lambda)\right)^{-1}. \quad (65)$$

4.2 Formal approximation of the first eigenvalue and corresponding eigenfunction

We look for an approximation $\lambda_N^{(0)}$ of the first eigenvalue of (54)–(57) and an approximation $\sigma_N(\mathbf{x}, \lambda_N^{(0)})$ of the corresponding eigenfunction. We use the representation

$$\sigma_N(\mathbf{x}, \lambda_N^{(0)}) = \sum_{j=1}^N C_j V^{(j)}(\mathbf{x}, \lambda_N^{(0)}), \quad (66)$$

with the coefficients C_j being unknown intensities of inertia forces exerted by the inclusions on the membrane. The approximation (66) satisfies (54) and the boundary condition (55) on $\partial\Omega$.

The boundary conditions (56), (57) yield the system of algebraic equations for the coefficients C_j .

4.2.1 The algebraic system

When $\mathbf{x} \in \partial F_\varepsilon^{(k)}$, we use (66), together with (56), (57), to deduce the following relations

$$-\frac{1}{\gamma_k \lambda_N^{(0)}} T^{(k)}[\sigma_N] = C_k + \sum_{j \neq k} C_j V^{(j)}(\mathbf{x}, \lambda_N^{(0)}), \quad \mathbf{x} \in \partial F_\varepsilon^{(k)}, \quad k = 1, \dots, N, \quad (67)$$

where

$$T^{(k)}[\sigma_N] = \int_{\partial F_\varepsilon} \frac{\partial \sigma_N}{\partial n} ds. \quad (68)$$

Taking into account that

$$T^{(k)}[\sigma_N] = -\beta_\varepsilon^{(k)} C_k + O(\varepsilon |\log \varepsilon|), \quad (69)$$

and expanding $V^{(j)}(\mathbf{x}, \lambda_N^{(0)})$ in the series form near $\mathbf{x} = \mathbf{O}^{(k)}$, $k \neq j$, we arrive at the algebraic system equations for the coefficients C_k

$$C_k \left(1 - \frac{\beta_\varepsilon^{(k)}}{\gamma_k \lambda_N^{(0)}}\right) + \sum_{j \neq k} C_j V^{(j)}(\mathbf{O}^{(k)}, \lambda_N^{(0)}) = 0, \quad k = 1, \dots, N, \quad (70)$$

and furthermore, taking into account (64) we approximate the coefficients C_j as non-trivial solutions of the following homogenous system

$$\{\mathbf{I} - \mathbf{D}_\varepsilon(\lambda_N^{(0)}) + \mathbf{S}_\varepsilon(\lambda_N^{(0)})\} \mathbf{C} = 0, \quad (71)$$

where $\mathbf{C} = (C_1, \dots, C_N)^T$, \mathbf{I} is the identity matrix, and $\mathbf{D}_\varepsilon(\lambda_N^{(0)})$, $\mathbf{S}_\varepsilon(\lambda_N^{(0)})$ are $N \times N$ matrices defined as

$$\begin{aligned} \mathbf{D}_\varepsilon(\lambda_N^{(0)}) &= \frac{1}{\lambda_N^{(0)}} \text{diag}\left\{\frac{\beta_\varepsilon^{(1)}}{\gamma_1}, \dots, \frac{\beta_\varepsilon^{(N)}}{\gamma_N}\right\}, \\ (\mathbf{S}_\varepsilon(\lambda_N^{(0)}))_{kj} &= \beta_\varepsilon^{(j)} G_\Omega(\mathbf{O}^{(k)}, \mathbf{O}^{(j)}, \lambda_N^{(0)})(1 - \delta_{jk}). \end{aligned} \quad (72)$$

The approximation $\lambda_N^{(0)}$ of the first eigenvalue is defined from the equation

$$\det\{\mathbf{I} - \mathbf{D}_\varepsilon(\lambda_N^{(0)}) + \mathbf{S}_\varepsilon(\lambda_N^{(0)})\} = 0. \quad (73)$$

Given $\lambda_N^{(0)}$, the approximation σ_N of the eigenfunction is defined by (66), with the coefficients C_j obtained from (71).

We also note that the above algorithm equally applies to the Dirichlet problem (54)–(56), with A_j being zero in the right-hand side of (56). In this case, the matrix term \mathbf{D}_ε in (71) is replaced by zero.

The asymptotic scheme, described here, does not apply when $\lambda_N^{(0)}$ is close to the elements of the discrete spectrum of the Dirichlet problem in Ω .

4.3 An example

The above scheme is applicable to sparse clusters or dense one-dimensional clusters of inertial inclusions, where the minimum distance between the inclusions within the cluster exceeds $d = O(\text{diam}(\Omega)/N)$. When the masses m_j of small inclusions increase, with the membrane mass density ρ remaining constant, coefficients γ_j in (57) also increase and the first eigenvalue of (54)–(57) decreases accordingly.

However, we obtain a different problem in the limit when the masses m_j tend to infinity, with ρ being finite. The corresponding formulation will be the Dirichlet eigenvalue problem (54)–(56), with the homogeneous boundary conditions (56). The equation (71), with \mathbf{D}_ε being zero, defines the approximation for the first eigenfrequency.

Here, we discuss an example, where the results of the asymptotic approximation are compared to the an alternative numerical simulation in COMSOL Multi-Physics.

When the domain Ω is the disk $\{\mathbf{x} : |\mathbf{x}| < R\}$ of radius R with the centre at the origin, and $\mathbf{y} \neq \mathbf{0}$, the regular part of Green's function in (60) and (65) is evaluated with the use of Graf's addition formula, as follows

$$R_\Omega(\mathbf{x}, \mathbf{y}, \lambda) = \text{Re}\left\{\sum_{n=-\infty}^{\infty} \alpha_n J_n(\sqrt{\lambda}|\mathbf{x}|) e^{in\theta_{x,0}}\right\}, \quad (74)$$




Figure 2: Asymptotic approximation of the eigenfunction corresponding to the first eigenvalue of the Dirichlet problem in the domain with the cluster of small inclusions placed along a contour. The approximation is $\lambda_N^{(0)} = 0.30678$. The main region is a disk of radius $R = 7$. Small circular rigid inclusions of radii $r = 0.2$ have their centres along the circle of radius 2.

where the coefficients α_n are given in the form

$$\alpha_n = \frac{iH_n^{(1)}(\sqrt{\lambda}R)J_n(\sqrt{\lambda}|\mathbf{y}|)}{4J_n(\sqrt{\lambda}R)}e^{-in(\theta_{0,y}+\pi)}, \quad (75)$$

where $\theta_{x,0}$ is the polar angle of \mathbf{x} with respect to the centre $\mathbf{0}$ of the disk, and $\theta_{0,y}$ is the polar angle of $\mathbf{0}$ with respect to \mathbf{y} .

When $\mathbf{y} = \mathbf{0}$, we have

$$R_\Omega(\mathbf{x}, \mathbf{0}, \lambda) = -\frac{Y_0(\sqrt{\lambda}R)}{4J_0(\sqrt{\lambda}R)}J_0(\sqrt{\lambda}|\mathbf{x}|). \quad (76)$$

Figure 2 shows the asymptotic approximation of the eigenfunction corresponding to the first eigenvalue of the Dirichlet problem in the domain Ω_N with the cluster of small inclusions placed along a contour. The approximation is $\lambda_N^{(0)} = 0.30678$. An independent numerical simulation produced in COMSOL Multiphysics gives the first eigenvalue of $\lambda = 0.30816$, with an excellent agreement observed in Fig. 1 and Fig. 2.

5 Concluding remarks

We have given an outline of problems, where meso-scale asymptotic approximations provide an analytical insight to analysis of time-harmonic wave

problems in two-dimensional domains with small inclusions arranged in a sparse cluster or a one-dimensional clusters placed along a curve.

We note that logarithmic asymptotics, required for analysis of Dirichlet eigenvalue problems in domains with small inclusions often lead to a constraint of exponentially small size of the inclusions. On the other hand, if the rigid small inclusions are considered as “movable”, the inertia of the inclusion is used as an additional control parameter in the meso-scale asymptotic approximation.

Three classes of formulations discussed here show several directions where asymptotics of time-harmonic waves prove to be useful. In particular, in Section 2, for a finite mass cluster of small inertial inclusions we have derived a transmission condition across an inertial structured interface. While in Section 3, the quasi-static Green’s function has been successfully employed, together with the relative capacitary potential, it is essential to take into account wave scattering and reflection and to engage Green’s function for the Helmholtz operator in the analysis of the eigenvalue problem for a domain with a lower-dimensional cluster of small inclusions in Section 4. As illustrated, the derived asymptotic formulae provide good analytical insight and are straightforward to use in practical examples and computations.

Acknowledgements

V.G.M. acknowledges that this publication has been prepared with the support of the “RUDN University Program 5-100”. A.B.M would like to thank the EPSRC (UK) for its support through the Programme Grant no. EP/L024926/1. M.J.N gratefully acknowledges the support of the EU H2020 grant MSCA-IF-2016-747334-CAT-FFLAP.

References

- [1] V. Maz’ya, A. Movchan (2010) Asymptotic treatment of perforated domains without homogenization, *Mathematische Nachrichten*, Vol. 283 (1), 104–125.
- [2] V. Maz’ya, A. Movchan, M. Nieves (2013) *Green’s Kernels and Meso-Scale Approximations in Perforated Domains*, *Lecture Notes in Mathematics 2077*, Springer, Heidelberg, New York, Dordrecht, London.
- [3] V. Maz’ya, A. Movchan, M. Nieves (2010) Green’s kernels for transmission problems in bodies with small inclusions. *Operator Theory and Its Applications*. Eds. M. Levitin, D. Vassiliev, American Mathematical Society Translations, Series 2, Vol. 231, 127–171.

- [4] V. Maz'ya, A. Movchan, M. Nieves (2011) Mesoscale asymptotic approximations to solutions of mixed boundary value problems in perforated domains. *SIAM Journal on Multiscale Modeling and Simulation*, 9 (1), 424–448.
- [5] V.A. Marchenko, E.Y. Khruslov (2006) *Homogenization of Partial Differential Equations*, Birkhäuser, Boston.
- [6] V. Maz'ya, A. Movchan, M. Nieves (2016) Eigenvalue problem in a solid with many inclusions: asymptotic analysis. *SIAM Journal on Multiscale Modeling and Simulation*, 15 (2), 1003–1047, DOI: 10.1137/16M1079348.
- [7] V. Maz'ya, S. Nazarov, B. Plamenevskii (2000) *Asymptotic Theory of Elliptic Boundary Value Problems in Singularly Perturbed Domains*, Vols. 1–2, Birkhäuser, Boston.
- [8] C. Linton (1998): The Green's function for the two-dimensional Helmholtz equation in periodic domains, *Journal of Engineering Mathematics* 33, no. 4, 377–401.
- [9] H. Kurkcu, N. Nigan and F. Reitich (2009): Integral representation of Green's function for a linear array of acoustic point sources, *Journal of Computational Physics* 230, no 8, 75–95.
- [10] L. L. Foldy (1945) The multiple scattering of waves. I. General theory of isotropic scattering by randomly distributed scatterers, *Phys. Rev.*, 67, 107–119.
- [11] C. M. Linton and P. A. Martin (2004) Semi-infinite arrays of isotropic point scatterers. A unified approach, *SIAM J. Appl. Math.*, 64, 1035–1056.
- [12] O. Schnitzer, R.V. Craster (2017) Bloch Waves in an Arbitrary Two-Dimensional Lattice of Subwavelength Dirichlet Scatterers, *SIAM J. Appl. Math.*, 77(6), 2119–2135.
- [13] J. Saranan, G. Vainikko (2002): *Periodic Integral and Pseudodifferential Equations with Numerical Approximation*, Springer Monographs in Mathematics, Springer-Verlag Berlin Heidelberg.